

Impact of habitat management of small game species on bird communities in French Mediterranean scrublands

N. Froustey^{1,2}, M. Guyot¹, M. Crétet¹, S. Aulagnier², L. Pelozuelo³, J. C. Ricci¹

Author affiliations:

¹ Institut Méditerranéen du Patrimoine Cynégétique et Faunistique, Vergèze, France
² INRAE, Université de Toulouse, Castanet-Tolosan, France
³ UMR 5245 CNRS-NPT-UPS, Toulouse, France

Corresponding author:

N Froustey
nicolas.froustey33@outlook.fr

Handling Editor:

Ian MacGregor-Fors

Received: 27/04/2023

Cond. acceptance: 18/10/2023

Final acceptance: 22/05/2024

Published: 30/05/2024

Abstract

Impact of habitat management of small game species on bird communities in French Mediterranean scrublands. In the French Mediterranean region, the progressive abandonment of traditional activities has led to the expansion of scrubland and landscape encroachment, thereby driving biodiversity loss. Hunters participate in habitat management mostly by opening scrubland to increase the habitat of small game species. The aim of our study was to evaluate the contribution of such management to the conservation of non-targeted bird species. In three 30-hectare managed (MA) and unmanaged (UMA) areas we monitored birds in spring using the progressive frequency sampling method (EFP). Species richness (a proxy of alpha diversity) and beta diversity indexes were estimated to compare bird communities in MA and UMA neighbouring areas. We also studied the presence of priority species and the response of bird species along a gradient of management. Our results showed that habitat management of small game species was associated with higher bird species richness, including open-habitat specialists. Most of the Mediterranean bird species detected were present in managed areas, with coexistence of bird species from open and closed environments. Hunters' management of scrubland environments can result in areas of high conservation value for non-targeted bird species. This study could support guidelines for opening scrubland in order to preserve bird diversity.

Key words: Avifauna, Scrubland opening, Species richness, Conservation, Southern France

Resumen

Impacto de la gestión del hábitat de especies de caza menor en las comunidades de aves de los matorrales mediterráneos franceses. En la región del Mediterráneo francés, la progresiva disminución de las actividades tradicionales ha dado lugar a la expansión de los matorrales, que han invadido el territorio, lo que induce cierta pérdida de biodiversidad. Los cazadores participan en la gestión del hábitat principalmente mediante la apertura de los matorrales para ampliar el hábitat de las especies de caza menor. El objetivo de nuestro estudio fue evaluar la contribución de dicha gestión a la conservación de especies de aves no objetivo. En tres áreas gestionadas (MA) y no gestionadas (UMA) de 30 hectáreas, las aves fueron monitoreadas en primavera mediante el método de muestreo de frecuencia progresiva (EFP). Se estimaron los índices de riqueza de especies (un indicador de la diversidad alfa) y de diversidad beta para comparar las comunidades de aves entre áreas gestionadas y no gestionadas vecinas. También se estudiaron la presencia de especies prioritarias para la conservación y la respuesta de las especies de aves a lo largo de un gradiente de gestión. Nuestros resultados mostraron que la gestión del hábitat de especies de caza menor se asoció con el aumento de la riqueza de especies de aves, en especial las especialistas en hábitats abiertos. La mayoría de las especies de aves mediterráneas detectadas estaba presente en zonas gestionadas, donde convivían especies de aves de ambientes abiertos y cerrados. La gestión de los ambientes de matorrales por parte de los cazadores puede dar lugar a áreas de alto valor de conservación para especies de aves no objetivo. Este estudio podría respaldar las pautas para la apertura de matorrales con el fin de conservar la diversidad de aves.

Palabras clave: Avifauna, Apertura de matorrales, Riqueza de especies, Conservación, Sur de Francia

Cite:

Froustey N, Guyot M, Crétet M, Aulagnier S, Pelozuelo L, Ricci JC, 2024. Impact of habitat management of small game species on bird communities in French Mediterranean scrublands. *Animal Biodiversity and Conservation* 47.1, 75-88. DOI: <https://doi.org/10.32800/abc.2024.47.0075>

© [2024] Copyright belongs to the authors, who license the journal *Animal Biodiversity and Conservation* to publish the paper under a Creative Commons Attribution 4.0 License.

ISSN: 1578-665 X
eISSN: 2014-928 X

Introduction

Contemporary Mediterranean land ecosystems are the heritage of a wide variety of land uses, particularly in southern France (Brotons et al 2006). These ecosystems constitute a 'biodiversity hotspot' whose preservation is a major conservation challenge (Myers et al 2000). However, in recent decades, the strong expansion of scrubland resulting from the progressive abandonment of traditional agricultural practices has changed these landscapes (Blondel and Médail 2009, Sirami et al 2010).

Scrubland expansion is widely recognized as a major driver of biodiversity loss in Mediterranean landscapes (Preiss et al 1997, Sirami et al 2008, Vimal et al 2017, Brotons et al 2018). It is one of the causes of the decline of some emblematic bird species, such as the ortolan bunting *Emberiza hortulana* (Fonderlick et al 2010). It is also a cause of the decline of some insects, such as many Orthoptera (Russo 2007), and also of small game species (Delibes-Mateos et al 2009, 2012) such as the red-legged partridge *Alectoris rufa*, the Iberian hare *Lepus granatensis*, and the European rabbit *Oryctolagus cuniculus* (Delibes-Mateos et al 2008, Ferreira et al 2014, Casas and Garcia 2022). Drastic land use changes from the 1950s to the 1990s affected the distribution and abundance of these species in the Mediterranean area of the Iberian Peninsula (Vargas et al 2007, Delibes-Mateos et al 2009, 2012). Birds are relevant indicators of ecological changes within landscapes (Blondel 1965, Russo 2007, Aebscher et al 2016). Scrubland expansion is highly detrimental to the numerous Mediterranean bird species that favour open garrigues and rocky habitats, particularly priority species for conservation which rely on open habitats and/or depend on human activities (Vimal et al 2017), such as the southern grey shrike *Lanius meridionalis* and the woodchat shrike *Lanius senator* (Sfougaris et al 2014, Moreno-Rueda et al 2016). Their presence is the result of long and constant rural uses despite strong variations in intensity and extent. Owing to the significant changes in Mediterranean landscapes, many species associated with agricultural ecosystems are currently in decline, such as, for example, the vulnerable European goldfinch *Carduelis carduelis* and the European serin *Serinus serinus* (IUCN France et al 2016). The significant loss of open landscape species is partly compensated by the expansion of forest species benefiting from the maturation of scrublands (Lindenmayer et al 2000, Laiolo et al 2004, Sirami et al 2007, Brotons et al 2018).

Hunters are concerned by these changes and share an interest in biodiversity with conservationists (Alphandéry and Fortier 2007, Casas et al 2009, Ballon et al 2012, Caro et al 2014, Créois et al 2020, Gortázar and Fernandez-de-Simon 2022). In France, hunters play an essential political, economic (Schultz et al 2003), social (Stedman and Heberlein 2001) and ecological role in rural areas (Loveridge et al 2007, Ginelli 2009, Gallo and Pejchar 2016). In order to favour the maintenance or increase of small game species populations they release individual partridges and hares, and manage suitable habitats for the target species (Mustin et al 2011). In the Mediterranean region they restore small open areas in scrublands, they fund the plantation of

crops for wildlife, they dig out water bodies, and they may also provide additional food seasonally (Borralho et al 1998, Ponce-Boutin et al 2004, Casas et al 2022). These habitat management practices often benefit non-target species (Gallo and Pejchar 2016, Mustin et al 2018). However, such practices are usually restricted to small experimental areas where ecological impacts on non-target species are rarely considered (Arroyo and Beja 2002, Sánchez-García et al 2024). Thus, the effect of scrubland opening by hunters on non-game wildlife in the French Mediterranean region has so far received little attention, despite its possible conservation value (Martinez-Padilla et al 2002). The aim of our study was to evaluate the impact of such habitat management on bird assemblages in three scrublands in southern France by comparing pairs of neighbouring managed (MA) and unmanaged (UMA) areas. We hypothesized that habitat management for small game species should benefit non-targeted bird species through (1) a higher richness and beta diversity, (2) an increased number of priority species for conservation, and (3) differences in occurrence within open habitat species.

Material and methods

Study sites

We sampled three sites in the French Mediterranean region, all with a landscape structure dominated by scrubland (fig. 1). These three sites have the characteristic local Mediterranean climate, namely, hot and dry summers (average summer temperature: 24.0–25.5 °C; average summer rainfall: 12–23 mm) and mild and wet winters (average winter temperature: 6.5–8.0 °C; average winter rainfall: 38–50 mm; 15–25 days of frost): (1) Bourg-Saint-Andéol (BSA, 44° 22' 24" N, 4° 38' 39" E, Ardèche, Auvergne-Rhône-Alpes) is located in the vicinity of the Ardèche canyon and the Rhône valley. This area is covered by vegetation varying from the meso-Mediterranean to lower supra-Mediterranean levels, including holm oak *Quercus ilex*, downy oak *Quercus pubescens* and boxwood coppice *Buxus sempervirens*; (2) Lançon-Provence (LP, 43° 35' 36" N, 5° 07' 43" E, Bouches du Rhône, Provence-Alpes-Côte-d'Azur) is located near the brackish water lagoon 'Etang de Berre' on the Fare Mountain range near the Dura valley. The xeric landscape is made up of scrubland with kermes oaks *Quercus coccifera* and rosemary bushes *Salvia rosmarinus*, Ibero-Mediterranean limestone cliffs, and rocky escarpments; and (3) Montpeyroux (MP, 43° 41' 46" N, 3° 30' 25" E, Hérault, Occitanie) is located at the foothills of the Cevennes, below Mont Saint-Baudille, at the extreme south-west of the southern slope of the Seranne mountain. It includes vast limestone hills, and is covered by scrubland with holm oak and xeric species of bushes. LP and MP are part of the Natura 2000 network.

Habitat management types

Each studied site included an area which was managed for small game species. Three main types of habitat management were identified (fig. 2):




Fig. 1. Location and landscape structure of the three study sites (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux), in southern France, including managed (MA) and unmanaged (UMA) areas for small game species, and the 108 bird count points. (Source: IGN, production IMPCF 2024).

Fig. 1. Ubicación y estructura del territorio de los tres sitios de estudio (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux) en el sur de Francia, incluidas las áreas gestionadas (MA) y no gestionadas (UMA) para especies de caza menor y los 108 puntos de conteo de aves. (Fuente: IGN, producción IMPCF 2024).

(1) Scrubland opening/clearance: corridors and open spaces of 20 to 100 hectares named alveolar openings, created within the scrubland using mechanical shredding. These openings are regularly created in autumn or winter, and some can evolve to dry grasslands after years.

(2) Crops for wildlife: small and closed natural areas of scrubland are converted or rehabilitated into game crops. Each plot is sown with subterranean clover *Trifolium subterraneum*, white lupin *Lupinus albus*, or orchard grass *Dactylis glomerata*, and cultivated without chemical input.

(3) Food and water supplementation: food hopper systems built from recycled materials and set on openings, corridors, or alongside crops. Each local hunting society maintains them on a daily basis (table 1s in supplementary material). Ponds or permanent water bodies are created to mitigate summer drought.

Vegetation maps of each studied area were produced using remote sensing technology. Eight variables were derived from remote sensing data to describe composition, vertical vegetation structure and heights opening alveolar, road or forest trail, crops, bare soil, low scrubs (< 1.50 m), high scrubs (> 1.50 m), trees, and herb layer in each MA or UMA. All steps were performed with QGIS 3.22.6 and the Orfeo ToolBox application (OTB 8.0.1) (Grizonnet et al 2017).

Description of bird assemblages

Bird assemblages were described using a fixed-radius count point method, the progressive frequency sampling (EFP) defined by Blondel (1975) to study the composition and structure of bird communities in various landscapes with minimal sampling effort (Archaux 2004). The radius, the time, the duration of the count points, and the distance between two neighbouring

points was further developed by other authors (Hutto et al 1986, Ralph et al 1995). This frequential sampling method enables the study of the composition and structure of bird communities in various landscapes.

Within each of the 30 ha MAs and UMAs spaced 1 to 1.5 km apart at each site according to Balent and Courtiade (1992), we established 18 fixed point-counts spaced 300 m apart to avoid overlap and ensure the independence of records. At each site and each day, we sampled the same number of point counts in MA and UMA. Bird species seen or heard, in flight or landed, were recorded within a 100-m radius surrounding the observer for 20 minutes, following Petit et al (1995) and Sirami et al (2007). Two Mediterranean bird experts conducted the field surveys simultaneously. They visited each MA and UMA several times, alternatively from 6:00 to 11:00 am between April 27 and June 3 in 2021 and 2022. Due to weather conditions, they sampled 42 point-counts per area at BSA, 48 point-counts per area at LP, and 57 point-counts per area at MP.

Species richness

Indices to estimate biodiversity are numerous but species richness is widely used in ecology (Scott et al 1987, Purvis and Hector 2000, Gotelli and Colwell 2001, Magurran 2021). We identified species richness as the number of species detected at a point count. We plotted the cumulative number of bird species as a function of the cumulative number of point count samples surveyed in MAs and UMAs and for each site using 'iNEXT' R package (Hsieh et al 2016), and contrasted species richness values using 84% confidence intervals (Payton et al 2003), after equalizing the sample sizes by extrapolation. An exhaustivity index of the species richness was calculated for MA and UMA at each study site using the Monod formula, adapted from the Michaelis-Menten formula (Lauga and Joachim 1987).

Beta diversity

Beta diversity evaluates the difference in species composition between two sites (Koleff et al 2003, Baselga 2010). The Sørensen index (Sørensen 1948) was used to compare species assemblages between MA and UMA at each site, and to check whether the recorded differences were due to additional species, or if areas were occupied by different communities. The values of the index vary from 0 (no similarity) to 1 (total similarity). The formula is as follows:

$$\beta_{sør} = 2 * A / 2 * A + X_{MA} + X_{UMA}$$

where A is the number of species recorded in both MAs and UMAs; X_{MA} is the number of species recorded only in the MA; X_{UMA} is the number of species recorded only in the UMA.

Priority species

We defined priority species according to their conservation and legal status (Cheylan 1995, Fonderlick 2007). These included species listed as vulnerable (VU) or endangered (EN) in the French IUCN Red list

(IUCN France et al 2016) and species listed in Annex I of the European Union Birds Directive (European Commission 2009) (table 2s in supplementary material). Species observed in flight and nocturnal birds were excluded from subsequent analyses because they cannot be considered direct users of the sampled area. We compared the prevalence (number of count points with record) of priority species between MAs and UMAs using chi-square-tests.

Effect of managed surface on bird ecological profile

It would be inadequate to evaluate the potential impact of MAs solely on the basis of species richness (Fusco et al 2021). It was thus important to consider the specific responses of the species according to their habitat preference listed in the *Inventaire National du Patrimoine Naturel* website (<https://inpn.mnhn.fr>). A species is considered generalist if it occurs in several types of habitats, and specialist if it occurs in only one type of habitat.

We therefore studied the occurrence of the 68 species recorded at the 147 point counts in MAs along a management gradient ranging from 0 to 49% of the surface. An occurrence frequency was calculated as the ratio between the number of point counts for which a species was present and the total number of point counts. Following Sirami (2006), we then considered a buffer zone of 100 m around each point count using remote sensing. Land use within each buffer was quantified in terms of managed surface and natural surface, before discretizing the managed surface into five classes that included the same amount of bird species data (0-4%, 4-13%, 13-18%, 18-27%, 27-49%).

Results

A total of 3,320 bird records were collected at the three sites over the two years of sampling. According to the study requirements, 2,322 records were kept, summarizing the occurrence of 71 bird species. The 998 records excluded from the analyses were: i) 958 records of flying birds, and ii), 40 records of landed raptors and nocturnal bird species.

Species richness and beta diversity

At the site level, species richness was 48 in BSA, 55 in LP and 55 in MP (fig. 3A). Species richness was significantly higher in MAs (68.00 ± 4.00 calculated species) than in UMAs (55.00 ± 3.15 calculated species; fig. 3B). This was confirmed at the study site level, with MA having significantly more bird species than UMA (BSA = MA: 49.16 ± 4.33 , UMA: 39.07 ± 3.27 calculated species; LP = MA: 55.81 ± 4.29 , UMA: 43.02 ± 3.73 calculated species; MP = MA: 52.5 ± 3 ; UMA: 43.39 ± 2.83 calculated species).

According to the accumulation curves and exhaustivity indices, around 90% of the estimated number of species were recorded in each area type of the three study sites (fig. 1s in supplementary material).

Sørensen indices showed that beta diversity was similar in the three study sites (table 1). However, the greater number of records in MAs was associated with a greater number of exclusive species (16 vs 3 in UMAs of the three sites).




Fig. 2. Drone shot illustrating a habitat managed for small game species in Bourg-Saint-Andéol ($44^{\circ} 22' 24''$ N, $4^{\circ} 38' 39''$ E). Each type of improvement is presented.

Fig. 2. Toma de dron que ilustra un hábitat gestionado para especies de caza menor en Bourg-Saint-Andéol ($44^{\circ} 22' 24''$ N, $4^{\circ} 38' 39''$ E). Se presentan todos los tipos de mejora de hábitat.

Priority species

Among the 71 bird species, 17 were considered priority species, either because of their French IUCN status (4 as 'Endangered', 9 as 'Vulnerable') or because they were listed in Annex I of the European Birds Directive (table 2). They included mainly the European turtle-dove *Streptopelia turtur*, the Dartford warbler *Sylvia*

undata and the European serin. Priority species were more prevalent in MA (fig. 4) and the difference was significant at BSA ($\chi^2 = 6.39$, $p < 0.05$), LP ($\chi^2 = 5.04$, $p < 0.05$) and MP ($\chi^2 = 10.6$, $p < 0.01$). On average, an observer had a 14.2% chance of observing a priority species when conducting a survey in MAs, compared to a 10.6% chance in UMA.




Fig. 3. A, bird species richness in managed (MA) and unmanaged (UMA) areas of the three study sites in southern France (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux). Each point corresponds to the number of species detected at a point count during a visit. Stars correspond to mean values. B, accumulation curves of bird species richness (y-axis) with 84 % confidence intervals estimated on 147 point counts (x-axis) in MAs and UMs using the R package 'iNEXT'.

Fig. 3. A, riqueza de especies de aves en áreas gestionadas (MA) y no gestionadas (UMA) de los tres sitios de estudio en el sur de Francia (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux). Cada punto corresponde al número de especies detectadas en un punto de conteo durante una visita. Las estrellas corresponden a valores medios. B, curvas de acumulación de riqueza de especies (eje y) con intervalos de confianza del 84 % estimados a partir de 147 puntos de conteo (eje x) en áreas gestionadas y no gestionadas utilizando el paquete "iNEXT" de R.

Ecological profiles of bird species

Sixty-eight species were grouped according to the average managed surface in the 100-m buffer zone of MA count points where they were detected (fig. 5). Three main ecological profiles were identified. Most species were present whatever the managed surface and can be considered generalist (45 species). These were priority species such as the European turtle-dove and the European greenfinch *Chloris chloris*, or common species such as the Eurasian magpie *Pica pica* and the common chaffinch *Fringilla coelebs*. The frequency of close habitat specialists (seven species), such as the endangered spectacled warbler *Sylvia conspicillata* and the Dartford

warbler, decreased with the managed surface of the scrubland in the buffer. In contrast, the open habitat specialists (16 species), such as the red-legged partridge *Alectoris rufa* and the common pheasant *Phasianus colchicus*, benefited from larger scrubland openings.

Discussion

Here we evaluated the outcome of small game habitat managed by hunters on non-targeted bird species in three French Mediterranean scrubland sites. With an overall occurrence of 71 species, the exhaustiveness of species richness close to 90% for each area type indicates that we recorded representative samplings

Table 1. Beta diversity indexes comparing bird communities recorded during count points in managed (MA) and unmanaged areas (UMA) for each study site of southern France: BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux.

Tabla 1. Índices de diversidad beta que permiten comparar las comunidades de aves registradas durante los puntos de conteo entre áreas gestionadas (MA) y no gestionadas (UMA) para cada sitio de estudio en el sur de Francia: BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux.

	Study sites					
	BSA		LP		MP	
Area	MA	UMA	MA	UMA	MA	UMA
Count points	42	42	48	48	57	57
Records	398	334	365	275	550	400
Species	46	37	53	41	52	43
Exclusive species	11	2	14	2	12	3
Sørensen index	0.843		0.830		0.842	

of the bird assemblages. Observability within the study sites was not an obstacle to detect the species in either MAs or UMs because 97% of our records were obtained by acoustic detection. Progressive frequency sampling (Blondel 1975) has been used previously in numerous studies in the Mediterranean region (e.g., Blondel et al 1978, Martin 1983, Fonderflick et al 2001). However, the relative abundance of each species was not estimated at each point count. This may be a limitation of our study in addition to the somewhat different management at the three sites (table 1s in supplementary material). Another recording method such as the index of point abundance may be required (Blondel et al 1970).

Effect of habitat management on bird richness and beta diversity

Our results support the former hypothesis, as habitats managed by hunters for small game species were more attractive for non-targeted bird species than unmanaged habitats, with both a higher species richness and a greater number of exclusive species. We observed a clear relationship between the diversity of bird species and the structural diversity of the sampled areas. Indeed, it has long been recognized that the structure of vegetation is the main ecological factor conditioning variability and differentiation of bird communities in the Mediterranean region (Prodon and Lebreton 1981). Nevertheless, our data lack an evaluation of bird species abundance in order to more accurately evaluate the influence of vegetation structure (Fonderflick 2007). The number of breeding pairs should be compared between MAs and UMs in a future study as suggested previously.

Within each study site, we were unable to identify the relative impact of each specific management practice on the non-targeted bird species at both local and landscape scales. We only considered scrubland opening for defining MAs, and punctual improvements such as feeders and trough were neglected. As no information was available about the area impacted by such improvements they could not be integrated

into the managed surface of the point count buffers. However, providing food and water to game species could also benefit non-targeted species (Gaudioso-Lacasa et al 2010, Arroyo et al 2013, Armenteros et al 2021), and even predators by increasing the number of prey (Beja et al 2009).

Benefit of habitat management on priority species

Our results also support the second hypothesis as we found MAs showed a positive effect on some priority species for conservation. This result is consistent with their selection of heterogeneous habitat at the landscape scale (Preiss et al 1997, Suárez-Seoane et al 2002). This is the case for threatened species inhabiting agroecosystems, such as the European goldfinch, the European serin, the European turtle-dove, and the European greenfinch, all of which were more frequently recorded in MAs than in UMs. Surprisingly, the endangered southern grey shrike, which also inhabits agroecosystems, did not benefit from habitat opening as did the open habitat specialist meadow pipit *Anthus pratensis*. On the other hand, the habitat management did not affect the occurrence of scrubland specialists such as the spectacled warbler or the Dartford warbler. This suggests that in the Mediterranean landscape, the role of habitat heterogeneity through MAs could be a key factor for conserving a number of threatened species.

Occurrence of open habitat specialists

Some of the bird species recorded in MAs do not belong to traditional Mediterranean bird communities of scrublands. This is due to a secondary colonization by species of open and human managed environments, including some priority species for conservation. This observation is consistent with previous studies carried out in the French Mediterranean region where most bird species with an unfavourable conservation status are linked to open and semi-open habitats (Fonderflick 2007, Vimal et al 2017, Brotons et al 2018). At a larger scale, in Europe, most priority species for conservation need a mixture of woodland,

Table 2. Occurrence of the 71 bird species recorded during count points in both the managed (MA) and unmanaged (UMA) area of the three study sites in southern France (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux). Species are listed according to their IUCN conservation status in France: EN, endangered; VU, vulnerable; NT, near threatened; LC, least concern (IUCN France et al 2016). Species whose IUCN status is followed by (*) are listed in Annex I of the European Bird Directive (European Commission 2009). Species flagged with (**) are not scrubland specialists.

Tabla 2. Avistamientos de las 71 especies de aves registradas durante los puntos de conteo tanto en las áreas gestionadas (MA) como en las no gestionadas (UMA) de los tres sitios de estudio en el sur de Francia (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux). Las especies se enumeran según el estado de conservación de la UICN en Francia: EN, en peligro; VU, vulnerable; NT, casi amenazado; LC, preocupación menor. Las especies cuyo estatus en la UICN va seguido de (*) se enumeran en el Anexo I de la Directiva Europea de Aves (Comisión Europea 2009). Las especies marcadas con (**) no son especialistas en matorrales.

Species	Valence	IUCN status	Number of occurrences					
			BSA		LP		MP	
			MA	UMA	MA	UMA	MA	UMA
<i>Acrocephalus melanopogon</i>	Specialist (**)	EN	0	0	1	0	0	0
<i>Lanius meridionalis</i>	Generalist	EN	0	0	1	2	2	2
<i>Sylvia conspicillata</i>	Specialist	EN	2	0	0	0	0	0
<i>Sylvia undata</i>	Specialist	EN	8	3	28	24	10	16
<i>Anthus pratensis</i>	Specialist	VU	0	0	1	0	0	1
<i>Carduelis carduelis</i>	Generalist	VU	3	0	6	1	4	1
<i>Chloris chloris</i>	Generalist	VU	3	1	0	0	3	2
<i>Lanius senator</i>	Generalist	VU	0	0	0	0	2	0
<i>Linaria cannabina</i>	Generalist	VU	1	0	1	0	3	3
<i>Pyrrhula pyrrhula</i>	Specialist (**)	VU	1	0	0	0	0	0
<i>Saxicola rubetra</i>	Specialist (**)	VU	0	0	1	0	0	0
<i>Serinus serinus</i>	Generalist	VU	0	0	3	1	26	6
<i>Streptopelia turtur</i>	Generalist	VU	27	20	13	4	36	15
<i>Coracias garrulus</i>	Generalist	NT (*)	0	0	1	1	0	0
<i>Lanius collurio</i>	Generalist	NT (*)	0	0	2	0	0	0
<i>Locustella naevia</i>	Generalist	NT	2	0	0	0	1	1
<i>Phylloscopus trochilus</i>	Generalist	NT	9	2	5	2	3	5
<i>Saxicola rubicola</i>	Generalist	NT	0	0	2	2	1	0
<i>Sylvia borin</i>	Generalist	NT	3	3	0	0	0	0
<i>Sylvia melanocephala</i>	Generalist	NT	26	25	45	40	48	53
<i>Acrocephalus scirpaceus</i>	Specialist (*)	LC	0	0	0	0	1	0
<i>Aegithalos caudatus</i>	Generalist (*)	LC	5	6	3	0	6	2
<i>Alectoris rufa</i>	Generalist	LC	10	3	13	14	22	6
<i>Anthus campestris</i>	Generalist	LC (*)	0	0	1	4	0	0
<i>Anthus trivialis</i>	Generalist	LC	0	0	0	0	1	0
<i>Caprimulgus europaeus</i>	Generalist	LC (*)	0	0	0	0	0	1
<i>Certhia brachydactyla</i>	Generalist	LC	3	4	2	0	5	2
<i>Certhia familiaris</i>	Specialist	LC	0	0	0	0	1	0
<i>Columba palumbus</i>	Generalist	LC	19	15	6	3	13	11
<i>Corvus corax</i>	Generalist	LC	3	0	0	0	0	0
<i>Corvus corone</i>	Generalist	LC	3	0	4	2	4	1
<i>Corvus monedula</i>	Generalist	LC	0	0	3	2	0	0
<i>Cuculus canorus</i>	Generalist	LC	1	3	4	1	18	11
<i>Cyanistes caeruleus</i>	Generalist	LC	9	16	8	3	8	6
<i>Dendrocopos major</i>	Specialist (**)	LC	1	0	0	0	0	0
<i>Emberiza calandra</i>	Specialist (**)	LC	2	0	3	2	6	0
<i>Emberiza cia</i>	Specialist (**)	LC	0	0	0	1	0	0
<i>Emberiza cirlus</i>	Generalist	LC	4	2	6	3	8	2
<i>Erythacus rubecula</i>	Generalist	LC	9	28	4	2	4	5
<i>Fringilla coelebs</i>	Generalist	LC	33	27	26	15	48	48
<i>Galerida cristata</i>	Generalist	LC	0	0	1	0	1	0
<i>Garrulus glandarius</i>	Generalist	LC	12	14	4	7	3	9

Table 2. (Cont.)

Species	Valence	IUCN status	Number of occurrences					
			BSA		LP		MP	
			MA	UMA	MA	UMA	MA	UMA
<i>Hippolais polyglotta</i>	Generalist	LC	3	0	6	5	12	8
<i>Lophophanes cristatus</i>	Generalist	LC	0	0	4	0	1	0
<i>Lulula arborea</i>	Generalist	LC	7	6	10	12	17	3
<i>Luscinia megarhynchos</i>	Generalist	LC	38	21	34	27	45	43
<i>Merops apiaster</i>	Generalist	LC	1	1	2	1	2	0
<i>Motacilla alba</i>	Generalist	LC	0	0	2	0	0	0
<i>Oriolus oriolus</i>	Generalist	LC	5	0	2	1	7	2
<i>Parus major</i>	Generalist	LC	19	20	12	9	21	14
<i>Passer domesticus</i>	Generalist	LC	0	0	0	0	1	0
<i>Phasianus colchicus</i>	Generalist	LC	2	2	21	21	3	3
<i>Phoenicurus ochruros</i>	Specialist (**)	LC	1	1	1	5	2	1
<i>Phoenicurus phoenicurus</i>	Specialist (**)	LC	12	2	7	4	15	11
<i>Phylloscopus borealis</i>	Generalist	LC	6	3	3	0	5	3
<i>Phylloscopus collybita</i>	Specialist	LC	1	2	1	1	3	2
<i>Pica pica</i>	Generalist	LC	1	3	8	13	2	3
<i>Picus viridis</i>	Generalist	LC	3	8	3	1	1	0
<i>Prunella modularis</i>	Specialist	LC	1	1	0	0	0	0
<i>Regulus ignicapilla</i>	Generalist	LC	8	2	2	0	10	3
<i>Sitta europaea</i>	Generalist	LC	0	0	0	0	3	0
<i>Streptopelia decaocto</i>	Generalist	LC	5	5	1	4	1	1
<i>Sturnus vulgaris</i>	Generalist	LC	4	1	1	0	0	0
<i>Sylvia atricapilla</i>	Specialist	LC	21	25	7	4	17	15
<i>Sylvia cantillans</i>	Specialist	LC	22	19	20	14	17	13
<i>Sylvia communis</i>	Specialist	LC	0	0	1	0	0	1
<i>Sylvia hortensis</i>	Generalist	LC	0	1	3	2	12	18
<i>Troglodytes troglodytes</i>	Specialist	LC	0	1	0	0	5	3
<i>Turdus merula</i>	Generalist	LC	35	34	6	8	28	36
<i>Turdus philomelos</i>	Generalist	LC	0	0	0	1	0	0
<i>Upupa epops</i>	Generalist	LC	4	4	10	6	32	8
Number of point counts			42	42	48	48	57	57




Fig. 4. Occurrences of priority species in count points surveyed in managed (MA) and unmanaged (UMA) areas of the three study sites in southern France: BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux. Stars indicate significant differences (chi-squared tests, $p < 0.05$).

Fig. 4. Avistamientos de especies prioritarias para la conservación en puntos de conteo estudiados en áreas gestionadas (MA) y no gestionadas (UMA) de los tres sitios de estudio en el sur de Francia: BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux. Las estrellas indican una diferencia significativa (pruebas chi-cuadrado, $p < 0,05$).

scrubland and rocky habitat within their foraging area (Tucker and Evans 1997). Nevertheless, most species exclusive to MAs are anthropophilic, with variable ecological amplitudes, and they occurred in managed habitats, either as to substitute or complement their preferred habitat. Described in a more general context, the sensitivity of bird species to environmental changes was shown to be linked to a sensory pattern whose main proxies are the scale of perception and the nature of the perceived elements (Sirami 2006). In our study, scrubland management has structured an agro-forestry bird assemblage in MAs. However, the ecological profile of most species living in the three scrublands identified them as generalists, known to be favoured by landscape opening (Dunning et al 1992, Brotons et al 2006, Caplat and Fonderlick 2009). Finally, we note that scrubland habitat management for small game species can fulfil biological requirements of many non-targeted bird species, and may be considered necessary to maintain habitat heterogeneity after agricultural abandonment.

Profit and loss of habitat management

Our results agree with findings in Spain (Caro et al 2014) and elsewhere in Europe (Preiss et al 1997, Stoate and Szczur 2001, Arroyo and Beja 2002, Stoate et al 2003, Sage et al 2005, Parish and Sotherton 2008, Santana et al 2017) where low-intensity management actions for small game had positive effects on non-targeted bird species. Several authors consider management practices that closely mimic ecological processes to be of added value for small game species (small agricultural plots, mosaic landscape or fire and mechanical shredding of scrubland cover) and have

positive effects on bird communities adapted to natural disturbance regimes (Smith et al 2001, Moreira et al 2001, Tapper 2005, Arroyo et al 2013, Aebischer et al 2016, Newey et al 2016). However, these studies have been conducted on farmlands and differ from our framework of analyses, making comparisons with our study difficult. Moreover, the effects of a particular type of management on non-targeted bird species are often paradoxical, inducing both positive and negative effects (Gallo and Pejchar 2016, Mustin et al 2018).

Although mechanical shredding may be important for fire prevention (Legrand et al 1994), its impact on Mediterranean vegetation and wildlife is not yet fully understood. The treatment periods could have a negative impact on the breeding success of ground-nesting birds, on the flowering cycle of plants, and subsequently on the abundance of invertebrates. Moreover, food and water supplementation can easily become ecological traps for some species users. However, the consequences on non-targeted bird species have yet to be fully explored.

Conclusion

Besides preventing fires, which are common in the region, the opening of scrublands appears to be of value for several bird species of high conservation priority.

In the present study, we were unable to identify an optimal threshold of management to support non-targeted bird species. Carrying out a study on a larger set of MAs for a longer period of time could provide more precise responses of non-targeted bird species to each management practice. Moreover, estimating the relative abundance of bird species in MAs based on




Fig. 5. Frequency of occurrence of the 68 bird species recorded in the 147 point counts classified in five equal frequency classes of managed surface in the three study sites in southern France. Species are ranked in ascending order according to the average of managed surface within a 100-m buffer around the count points where they were detected.

Fig. 5. Frecuencia de avistamientos de las 68 especies de aves registradas en los 147 puntos de conteo clasificada en cinco clases iguales de frecuencia de superficie gestionada en los tres sitios de estudio en el sur de Francia. Las especies están ordenadas de menor a mayor según el promedio de superficie gestionada dentro de los 100 metros alrededor de los puntos de conteo en los que fueran detectadas.

breeding pair census would further support the role of MAs for the conservation of non-targeted bird species. Nevertheless, until such results are available, habitat management of small game species in scrublands by hunters at a local scale can benefit the conservation of several bird species. Agricultural or forest management, including food and water provision, should be a further step (Sánchez-García et al 2024).

References

- Aebischer NJ, Bailey CM, Gibbons DW, Morris AJ, Peach WJ, Stoate C, 2016. Twenty years of local farmland bird conservation: the effects of management on avian abundance at two UK demonstration sites. *Bird Study* 63(1), 10-30. DOI: [10.1080/00063657.2015.1090391](https://doi.org/10.1080/00063657.2015.1090391)
- Alphandéry P, Fortier A, 2007. A new approach to wildlife management in France: regional guidelines as tools for the conservation of biodiversity. *Sociología Ruralis* 47(1), 42-62. DOI: [10.1111/j.1467-9523.2007.00426.x](https://doi.org/10.1111/j.1467-9523.2007.00426.x)
- Archaux F, 2004. Breeding upwards when climate is becoming warmer: no bird response in the French Alps. *Ibis* 146(1), 138-144. DOI: [10.1111/j.1474-919X.2004.00246.x](https://doi.org/10.1111/j.1474-919X.2004.00246.x)
- Armenteros JA, Caro J, Sanchez-Garcia C, Arroyo B, Perez JA, Gaudioso VR, Tizado EJ, 2021. Do non-target species visit feeders and water troughs targeting small game? A study from farmland Spain using camera-trapping. *Integrative Zoology* 16(2), 226-239. DOI: [10.1111/1749-4877.1249](https://doi.org/10.1111/1749-4877.1249)
- Arroyo B, Beja P, 2002. *Impact of hunting management practices on biodiversity*. Report Workpackage 2 to REGHAB Project, European Commission, Brussels. Available online at <http://digital.csic.es/bitstream/10261/8260/1/WP2-repport.pdf>
- Arroyo B, Delibes-Mateos M, Caro J, Estrada A, Mougeot F, Diaz-Fernández S, Casas F, Viñuela J, 2013. Efecto de la gestión para las especies de caza menor sobre la fauna no cinegética. *Ecosistemas* 22(2), 27-32. Available online at <https://revistaecosistemas.net/index.php/ecosistemas/article/view/734>
- Balent G, Courtiade B, 1992. Modelling bird communities/landscape patterns relationships in a rural area of South-Western France. *Landscape Ecology* 6, 195-211. DOI: [10.1007/BF00130031](https://doi.org/10.1007/BF00130031)
- Ballon P, Ginelli L, Vollet D, 2012. Les services rendus par la chasse en France : regards croisés en écologie, économie et sociologie. *Revue forestière française* 64(3), 305-318. DOI: [10.4267/2042/48439](https://doi.org/10.4267/2042/48439)
- Baselga A, 2010. Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography* 19(1), 134-143. DOI: [10.1111/j.1466-8238.2009.00490.x](https://doi.org/10.1111/j.1466-8238.2009.00490.x)
- Beja P, Gordinho L, Reino L, Loureiro F, Santos-Reis M, Borrallo R, 2009. Predator abundance in relation to small game management in southern Portugal: conservation implications. *European Journal of Wildlife Research* 55, 227-238. DOI: [10.1007/s10344-008-0236-1](https://doi.org/10.1007/s10344-008-0236-1)
- Blondel J, 1965. Etude des populations d'oiseaux dans une garrigue méditerranéenne : description du milieu, de la méthode de travail et exposé des premiers résultats obtenus à la période de reproduction. *Revue d'Écologie (La Terre et La Vie)* 19(4), 311-341. DOI: [10.3406/revec.1965.4463](https://doi.org/10.3406/revec.1965.4463)
- Blondel J, 1975. L'analyse des peuplements d'oiseaux, éléments d'un diagnostic écologique. I. la méthode des échantillonnages fréquentiels progressifs (EFP). *Revue d'Écologie (La Terre et La Vie)* 29(4), 533-589. DOI: [10.3406/revec.1975.4903](https://doi.org/10.3406/revec.1975.4903)
- Blondel J, David P, Lepart J, Romane F, 1978. L'avifaune du mont Ventoux essai de synthèse biogéographique et écologique. *Revue d'Écologie (La Terre et La Vie)* Suppl. 1, 111-145, www.persee.fr/doc/revec_2022-0790_1978_sup_32_1_5154
- Blondel J, Ferry C, Frochot B, 1970. La méthode des indices ponctuels d'abondance (IPA) ou des relevés d'avifaune par "stations d'écoute". *Alauda* 38(1), 55-71.
- Blondel J, Médail F, 2009. The physical geography of the Mediterranean. Biodiversity and conservation. In: *The physical Geography of the Mediterranean*: 615-650 (JC Woodward, Ed). Oxford University Press, Oxford.
- Borrallo R, Rito A, Rego F, Simoes H, Pinto PV, 1998. Summer distribution of red-legged partridges *Alectoris rufa* in relation to water availability on Mediterranean farmland. *Ibis* 140(4), 620-625. DOI: [10.1111/j.1474-919X.1998.tb04707.x](https://doi.org/10.1111/j.1474-919X.1998.tb04707.x)
- Brotons L, Herrando S, Martin JL, 2006. Bird assemblages in forest fragments within Mediterranean mosaics created by wildfires. *Landscape Ecology* 19(6), 663-675. DOI: [10.1007/s10980-005-0165-2](https://doi.org/10.1007/s10980-005-0165-2)
- Brotons L, Herrando S, Sirami C, Katí V, Díaz M, 2018. Mediterranean forest bird communities and the role of landscape heterogeneity in space and time. In: *Ecology and Conservation of Forest Birds*: 318-349 (G Mikusinski, JM Roberge, RJ Fuller, Eds). Cambridge University Press, Cambridge.
- Caplat P, Fonderlick J, 2009. Area mediated shifts in bird community composition: a study on a fragmented Mediterranean grassland. *Biodiversity and Conservation* 18(11), 2979-2995. DOI: [10.1007/s10531-009-9620-8](https://doi.org/10.1007/s10531-009-9620-8)
- Caro J, Delibes-Mateos M, Estrada A, Borrallo R, Gordinho L, Reino L, Beja P, Arroyo B, 2014. Effects of hunting management on Mediterranean farmland birds. *Bird Conservation International* 25(2), 166-181. DOI: [10.1007/S0959270914000197](https://doi.org/10.1007/S0959270914000197)
- Casas F, Duarte J, Gonçalves D, Meriggi A, Morales MB, Ponce F, Reino L, Traba J, Vargas JM, 2022. Habitat use and selection: is habitat management the key to restore red-legged partridge populations? In: *The future of the red-legged partridge*. *Wildlife research monographs*: 45-67 (F Casas, JT Garcia, Eds). Cham, Springer. DOI: [10.1007/978-3-030-96341-5_3](https://doi.org/10.1007/978-3-030-96341-5_3)
- Casas F, Garcia JT, 2022. *The Future of the Red-legged Partridge: Science, Hunting and Conservation*, Vol. 6. Springer Nature, Ciudad Real.
- Casas F, Mougeot F, Viñuela J, Bretagnolle V, 2009. Effects of hunting on the behaviour and spatial distribution of farmland birds: importance of hunting-free refuges in agricultural areas. *Animal Conservation* 12(4), 346-354. DOI: [10.1111/j.1469-1795.2009.00259.x](https://doi.org/10.1111/j.1469-1795.2009.00259.x)
- Cheylan M, 1995. Les Reptiles du Paléarctique occidental : diversité et conservation. Doctoral dissertation, Paris, EPHE. Available online at <https://www.theses.fr/1995EPHE3034>
- Crétois B, Linnell JD, Grainger M, Nilsen EB, Rød JK, 2020. Hunters as citizen scientists: contributions to biodiversity monitoring in Europe. *Global Ecology and Conservation* 23: e01077. DOI: [10.1016/j.gecco.2020.e01077](https://doi.org/10.1016/j.gecco.2020.e01077)
- Delibes-Mateos M, Farfán MA, Olivero JA, Márquez L, Vargas JM, 2009. Long-term changes in game species over a long period of transformation in the Iberian Mediterranean landscape. *Environmental Management* 43(6), 1256-1268. DOI: [10.1007/s00267-009-9297-5](https://doi.org/10.1007/s00267-009-9297-5)
- Delibes-Mateos M, Farfan MA, Olivero J, Vargas JM, 2012. Impact of land-use changes on red-legged partridge conservation in the Iberian Peninsula. *Environmental Conservation* 39(4), 337-346. DOI: [10.1017/S0376892912000100](https://doi.org/10.1017/S0376892912000100)
- Delibes-Mateos M, Ferreras P, Villafuerte R, 2008. Key role of European rabbits in the conservation of the Western Mediterranean basin hotspot. *Conservation Biology* 22(5), 1106-1117. DOI: [10.1111/j.1523-1739.2008.00993.x](https://doi.org/10.1111/j.1523-1739.2008.00993.x)
- Dunning JB, Danielson JB, Pulliam HR, 1992. Ecological processes that affect populations in complex landscapes. *Oikos* 65(1), 169-175. DOI: [10.2307/10.2307/3544901](https://doi.org/10.2307/10.2307/3544901)
- European Commission, 2009. Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the conservation of wild birds. *Official Journal of the European Union* 20, 7-25.
- Ferreira C, Touza J, Rouco C, Díaz-Ruiz F, Fernández de Simón J, Ríos Saldaña, Ferreras P, Villafuerte R, Delibes-Mateos M, 2014. Habitat management as a generalized tool to boost European rabbit *Oryctolagus cuniculus* populations in the Iberian Peninsula: a cost-effectiveness analysis. *Mammal Review* 44(1), 30-43. DOI: [10.1111/mam.12006](https://doi.org/10.1111/mam.12006)
- Fonderlick J, 2007. Conséquences de la fermeture et de la fragmentation des milieux ouverts sur l'avifaune nicheuse des Causses. Diplôme EPHE, Montpellier. Available online at <https://www.researchgate.net/publication/278302627>
- Fonderlick J, Caplat P, Lovaty F, Thévenot M, Prodon R, 2010. Avifauna trends following changes in a Mediterranean upland pastoral system. *Agriculture, Ecosystems and Environment* 137(3-4), 337-347. DOI: [10.1016/j.agee.2010.03.004](https://doi.org/10.1016/j.agee.2010.03.004)
- Fonderlick J, Thevenot M, Destre R, 2001. Le peuplement d'oiseaux du Causse Méjean (Lozère, France): état actuel, évolution historique et perspectives d'avenir. *Revue d'Écologie (La Terre et La Vie)* 56(2), 173-192. DOI: [10.3406/revec.2001.2357](https://doi.org/10.3406/revec.2001.2357)
- Fusco J, Walker E, Papaix J, Debolini M, Bondeau A, Barnagaud JY, 2021. Land use changes threaten bird taxonomic and functional diversity across the Mediterranean Basin: a spatial analysis to prioritize monitoring for conservation. *Frontiers in Ecology and Evolution* 9, 612356. DOI: [10.3389/fevo.2021.612356](https://doi.org/10.3389/fevo.2021.612356)
- Gallo T, Pejchar L, 2016. Improving habitat for game animals has mixed consequences for biodiversity conservation. *Biological Conservation* 197, 47-52. DOI: [10.1016/j.biocon.2016.02.032](https://doi.org/10.1016/j.biocon.2016.02.032)
- Gaudioso Lacasa VR, Sánchez García-Abad C, Prieto Martín R, Bartolomé Rodríguez DJ, Pérez Garrido JA, Alonso de La Varga ME, 2010. Small game water troughs in a Spanish agrarian pseudo steppe: visits and water site choice by wild fauna. *European Journal of Wildlife Research* 56, 591-599. DOI: [10.1007/s10344-009-0352-6](https://doi.org/10.1007/s10344-009-0352-6)

- Ginelli L, 2009. Chasse-gestion, chasse écologique, chasse durable. Enjeux autour de l'écolégisation d'une pratique en crise. In: *Chasse, Territoires et Développement durable. Outils d'analyse, enjeux et perspectives. Économie rurale* 327-328(1-2), 38-51. DOI: [10.4000/economierurale.3342](https://doi.org/10.4000/economierurale.3342)
- Gortázar C, Fernandez-de-Simon J, 2022. One tool in the box: the role of hunters in mitigating the damages associated to abundant wildlife. *European Journal of Wildlife Research* 68(3), 8-14. DOI: [10.1007/s10344-022-01578-7](https://doi.org/10.1007/s10344-022-01578-7)
- Gotelli NJ, Colwell RK, 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* 4(4), 379-391. DOI: [10.1046/j.1461-0248.2001.00230.x](https://doi.org/10.1046/j.1461-0248.2001.00230.x)
- Grizonnet M, Michel J, Poughon V, Ingla J, Savinaud M, Cresson R, 2017. Orfeo ToolBox: open source processing of remote sensing images. *Open Geospatial Data, Software and Standards* 2(1), 15. DOI: [10.1186/s40965-017-0031-6](https://doi.org/10.1186/s40965-017-0031-6)
- Hsieh TC, Ma K, Chao A, 2016. iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7(12), 1451-1456. DOI: [10.1111/210X.12613](https://doi.org/10.1111/210X.12613)
- Hutto RL, Pletschet SM, Hendricks P, 1986. A fixed-radius point count method for nonbreeding and breeding season use. *The Auk* 103(3), 593-602.
- Koleff P, Gaston KJ, Lennon JJ, 2003. Measuring beta diversity for presence-absence data. *Journal of Animal Ecology* 72(3), 367-382. DOI: [10.1046/j.1365-2656.2003.00710.x](https://doi.org/10.1046/j.1365-2656.2003.00710.x)
- Laiolo P, Dondero F, Ciliento E, Rolando A, 2004. Consequences of pastoral abandonment for the structure and diversity of the alpine avifauna. *Journal of Applied Ecology* 41(2), 294-304. DOI: [10.1111/j.0021-8901.2004.00893.x](https://doi.org/10.1111/j.0021-8901.2004.00893.x)
- Lauga J, Joachim J, 1987. Sampling bird communities by the E.F.P. method: mathematical study of the cumulative curve. *Acta Oecologica* 8(2), 117-124. DOI: <https://www.researchgate.net/publication/279542138>
- Legrand C, Etienne M, Rigolot E, 1994. Une méthode d'aide au choix des combinaisons techniques pour l'entretien des coupures de combustible. *Forêt méditerranéenne* 15(4), 397-408.
- Lindenmayer DB, Margules CR, Botkin DB, 2000. Indicators of biodiversity for ecologically sustainable forest management. *Conservation Biology* 14(4), 941-950. DOI: [10.1046/j.1523-1739.2000.98533.x](https://doi.org/10.1046/j.1523-1739.2000.98533.x)
- Loveridge AJ, Reynolds JC, Milner-Gulland EJ, 2007. Does sport hunting benefit conservation? In: *Key Topics in Conservation Biology*: 224-241 (DW Macdonald, K Service, Eds). Oxford University Press, Oxford.
- Magurran AE, 2021. Measuring biological diversity. *Current Biology* 31(19), R1174-R1177. DOI: [10.1016/j.cub.2021.07.049](https://doi.org/10.1016/j.cub.2021.07.049)
- Martin JL, 1983. The diagnose of density changes in island bird communities by the mean of point counts (progressive frequential sampling, E. F. P.). *Acta Oecologica, Ecologia Generalis* 4(2), 167-179.
- Martínez-Padilla J, Viñuela J, Villafuerte R, 2002. Socio-economic aspects of gamebird hunting, hunting bags, and assessment of the status of gamebird populations in REGHAB countries. Part 1: Socio-economics and cultural aspects of gamebird hunting. Instituto de Investigación en Recursos Cinegéticos, Ciudad Real. Available online at <http://hdl.handle.net/10261/8259>
- Moreira F, Ferreira PG, Rego FC, Bunting S, 2001. Landscape changes and breeding bird assemblages in northwestern Portugal: the role of fire. *Landscape Ecology* 16, 175-187. DOI: [10.1023/A:1011169614489](https://doi.org/10.1023/A:1011169614489)
- Moreno-Rueda G, Abril-Colón I, López-Orta A, Álvarez-Benito I, Castillo-Gómez C, Comas M, Rivas JM, 2016. Breeding ecology of the southern shrike (*Lanius meridionalis*) in an agrosystem of south-eastern Spain: the surprisingly excellent breeding success in a declining population. *Animal Biodiversity and Conservation* 39(1), 89-98. DOI: [10.32800/abc.2016.39.0089](https://doi.org/10.32800/abc.2016.39.0089)
- Mustin K, Arroyo B, Beja P, Newey S, Irvine RJ, Kestler J, Redpath SM, 2018. Consequences of game bird management for non-game species in Europe. *Journal of Applied Ecology* 55(5), 2285-2295. DOI: [10.1111/1365-2664.13131](https://doi.org/10.1111/1365-2664.13131)
- Mustin K, Newey S, Irvine J, Arroyo B, Redpath S, 2011. Biodiversity impacts of game bird hunting and associated management practices in Europe and North America. RSPB Report, Available at https://fp7hunt.net/Portals/HUNT/Reports/RSPB_ReportFINAL_Covers.pdf
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J, 2000. Biodiversity hotspots for conservation priorities. *Nature* 403(6772), 853-858. DOI: [10.1038/35002501](https://doi.org/10.1038/35002501)
- Newey S, Mustin K, Bryce R, Fielding D, Redpath S, Bunnefeld N, Bronwen D, Irvine RJ, 2016. Impact of management on avian communities in the Scottish Highlands. *Plos One* 11, e0155473. DOI: [10.1371/journal.pone.0155473](https://doi.org/10.1371/journal.pone.0155473)
- Parish DM, Sotherton NW, 2008. Landscape-dependent use of a seed-rich habitat by farmland passerines: relative importance of game cover crops in a grassland versus an arable region of Scotland. *Bird Study*, 55(1), 118-123. DOI: [10.1080/00063650809461512](https://doi.org/10.1080/00063650809461512)
- Payton ME, Greenstone MH, Schenker N, 2003. Overlapping confidence intervals or standard error intervals: what do they mean in terms of statistical significance? *Journal of Insect Science* 3(1), 34-40. DOI: [10.1093/jis/3.1.34](https://doi.org/10.1093/jis/3.1.34)
- Petit DR, Petit LJ, Saab VA, Martin TE, 1995. Fixed-radius point counts in forests: factors influencing effectiveness and efficiency. In: *Monitoring Bird Populations by Point Counts*: 49-56 (CJ Ralph, JR Sauer, S Droege, Eds). USDA Forest Service General Technical Report PSW-GTR-149.
- Ponce-Boutin F, Mathon JF, Le Brun T, Moutarde C, Corda E, Kmiec L, 2004. Aménagements des milieux et perdrix rouge en collines méditerranéennes françaises. *Faune Sauvage* 262, 42-46.
- Preiss E, Martin JL, Debussche M, 1997. Rural depopulation and recent landscape changes in a Mediterranean region: consequences to the breeding avifauna. *Landscape Ecology* 12(1), 51-61. DOI: [10.1007/BF02698207](https://doi.org/10.1007/BF02698207)
- Prodon R, Lebreton JD, 1981. Breeding avifauna of a Mediterranean succession: the holm oak and cork oak series in the eastern Pyrenees. 1. Analysis and modelling of the structure gradient. *Oikos* 37(1), 21-38. DOI: [10.2307/3544069](https://doi.org/10.2307/3544069)
- Purvis A, Hector A, 2000. Getting the measure of biodiversity. *Nature* 405(6783), 212-219. DOI: [10.1038/35012221](https://doi.org/10.1038/35012221)
- Ralph CJ, Droege S, Sauer JR, 1995. Managing and monitoring birds using point counts: standards and applications. In: *Monitoring Bird Populations by Point Counts*: 161-168 (CJ Ralph, JR Sauer, S Droege, Eds). USDA Forest Service General Technical Report PSW-GTR-149.
- Russo D, 2007. Effects of land abandonment on animal species in Europe: conservation and management implications. Integrated assessment of vulnerable ecosystems under global change in the European Union. Project report. European Commission, Community Research, Sustainable development, global change and ecosystems, Portici. Available online at <https://www.pik-potsdam.de/ateam/avec/paperrusso.pdf>
- Sage RB, Parish DM, Woodburn MI, Thompson PG, 2005. Songbirds using crops planted on farmland as cover for game birds. *European Journal of Wildlife Research* 51, 248-253. DOI: [10.1007/s10344-005-0114-z](https://doi.org/10.1007/s10344-005-0114-z)
- Sánchez-García C, Powolny T, Lormée H, Dias S, Sardà-Palomera F, Bota G, Arroyo B, 2024. Habitat management carried out by hunters in the European Turtle dove western flyway: opportunities and pitfalls for linking with sustainable hunting. *Journal for Nature Conservation* 78(1), 126561. DOI: [10.1016/j.jnc.2024.126561](https://doi.org/10.1016/j.jnc.2024.126561)
- Santana J, Reino L, Stoate C, Moreira F, Ribeiro PF, Santos JL, Rotenberry JT, Beja P, 2017. Combined effects of landscape composition and heterogeneity on farmland avian diversity. *Ecology and Evolution* 7(4), 1212-1223. DOI: [10.1002/ece3.2693](https://doi.org/10.1002/ece3.2693)
- Schulz JH, Millsbaugh JJ, Zekor DT, Washburn BE, 2003. Enhancing sport-hunting opportunities for urbanites. *Wildlife Society Bulletin* 31(2), 565-573, <http://www.jstor.org/stable/3784340>
- Scott JM, Csuti B, Jacobi JD, Estes JE, 1987. Species richness. *BioScience* 37(11), 782-788. DOI: [10.2307/1310544](https://doi.org/10.2307/1310544)
- Sfougaris AI, Plexida SG, Solomou AD, 2014. Assessing the effects of environmental factors on the presence and density of three shrike species in a continental and a coastal area of central Greece. *North-Western Journal of Zoology* 10(1), 101-109.
- Sirami C, 2006. Abandon des terres et avifaune: dynamiques spatiales et temporelles d'un paysage méditerranéen, Doctoral dissertation, Montpellier, École nationale supérieure agronomique. Available online at <https://www.theses.fr/2006ENSA0018>
- Sirami C, Brotons L, Burfield I, Fonderlick J, Martin JL, 2008. Is land abandonment having an impact on biodiversity? A meta-analytical approach to bird distribution changes in the north-western Mediterranean. *Biological Conservation* 141(2), 450-459. DOI: [10.1016/j.biocon.2007.10.015](https://doi.org/10.1016/j.biocon.2007.10.015)
- Sirami C, Brotons L, Martin JL, 2007. Vegetation and songbird response to land abandonment: from landscape to census plot. *Diversity and Distributions* 13(1), 42-52. DOI: [10.1111/j.1472-4642.2006.00297.x](https://doi.org/10.1111/j.1472-4642.2006.00297.x)
- Sirami C, Nesporous A, Cheylan JP, Marty P, Hvenegaard GT, Geniez P, Martin JL, 2010. Long-term anthropogenic and ecological dynamics of a Mediterranean landscape: impacts on multiple taxa. *Paysage et Urbanisme* 96(4), 214-223. DOI: [10.1016/j.landurbplan.2010.03.007](https://doi.org/10.1016/j.landurbplan.2010.03.007)
- Smith AA, Redpath SM, Campbell ST, Thirgood SJ, 2001. Meadow pipits, red grouse and the habitat characteristics of managed grouse moors. *Journal of Applied Ecology* 38(2), 390-400. DOI:

- 10.1046/j.1365-2664.2001.00601.x
- Sørensen TA, 1948. A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons. *Biologiske Skrifter, Kongelige Danske Videnskabernes Selskab* 5(4), 1-34. Available online at https://www.royalacademy.dk/Publications/High/2955_%C3%BBrensen,%20T%20horvald.pdf
- Stedman RC, Heberlein TA, 2001. Hunting and rural socialization: contingent effects of the rural setting on hunting participation. *Rural Sociology* 66(4), 599-617. DOI: 10.1111/j.1549-0831.2001.tb00086.x
- Stoate C, Szczur J, 2001. Could game management have a role in the conservation of farmland passerines? A case study from a Leicestershire farm. *Bird Study* 48(3), 279-292. DOI: 10.1080/00063650109461228
- Stoate C, Szczur J, Aebscher NJ, 2003. Winter use of wild bird cover crops by passerines on farmland in northeast England. *Bird Study* 50(1), 15-21. DOI: 10.1080/00063650309461285
- Suárez-Seoane S, Osborne PE, Baudry J, 2002. Responses of birds of different biogeographic origins and habitat requirements to agricultural land abandonment in northern Spain. *Biological Conservation* 105(3), 333-344. DOI: 10.1016/S0006-3207(01)00213-0
- Tapper SC, 2005. *Nature's gain: how gamebird management has influenced wildlife conservation*. The Game Conservancy Trust, Fordingbridge, Hampshire. Available online at <https://www.gwct.org.uk/media/249272/naturesgain.pdf>
- Tucker GM, Evans MI, 1997. *Habitats for Birds in Europe: a conservation strategy for the wider environment*. BirdLife International, Birdlife Conservation Series n°6), Cambridge.
- UICN France, MNHN, LPO, SEOF, ONCFS, 2016. *La Liste Rouge des espèces menacées en France : Oiseaux de France métropolitaine*. Available online at: <https://bit.ly/2zg7skP>
- Vargas JM, Farfan MA, Guerrero JC, Barbosa AM, Real R, 2007. Geographical and environmental correlates of big and small game in Andalusia (southern Spain). *Wildlife Research* 34(6), 498-506. DOI: 10.1071/WR06012
- Vimal R, Fonderlick J, Thompson JD, Pluvinet P, Debussche M, Cheylan M, Geniez P, Mathevet R, Lepart J, 2017. Integrating habitat diversity into species conservation in the Mediterranean mosaic landscape. *Basic and Applied Ecology* 22, 36-43. DOI: 10.1016/j.baae.2017.07.001

Acknowledgements

This research was conducted by the Institut Méditerranéen du Patrimoine Cynégétique et Faunistique. We thank all master's degree students: E Balandraud, F Moreau, M Cretet and ornithologists, especially P Dahler and Q Gaule, who collected data in the field. We thank the two anonymous reviewers and the associate editor for their comments that contributed to the improvement of this manuscript.

Author contributions

N Froustey, JC Ricci, L Pelozuelo and S Aulagnier conceived the original idea of the presented study. **N Froustey, Q Gaulé and P Dahler** conducted the fieldwork. **N Froustey, M Guyot and M Crétet** conducted the data analysis. **N Froustey and S Aulagnier** wrote the first draft of this publication. All these authors provided critical feedback and helped to shape the research, analysis and approved the final manuscript.

Conflicts of interest

No conflicts declared

Funding

The authors thank all financial contributors, especially Hunters Federal Union at the national (FNC), regional (FRC-PACA), and department levels: FDC 04-05-06-13-83-84 in Provence Alpes Côte d'Azur, FDC 2A-2B in Corsica, FDC 11-34-48-66-12 and 81 in Occitania, FDC 43-07-26 in Auvergne-Rhône-Alpes, and Regional Council Sud Provence Alpes Côte d'Azur.

Complete affiliations

N Froustey, M Guyot, M Crétet, JC Ricci, Institut Méditerranéen du Patrimoine Cynégétique et Faunistique, Site Expérimental Agri-Environnement les Bouillens, 30310 Vergèze, France

N Froustey, S Aulagnier, Comportement et Ecologie de la Faune Sauvage, INRAE, Université de Toulouse, 31326 Castanet-Tolosan, France
L Pelozuelo, Laboratoire d'écologie fonctionnelle et environnement, UMR 5245 CNRS-INPT, Université Toulouse 3 Paul Sabatier, Toulouse, France

Supplementary material

Table 1s. Habitat management made by hunters at the three study sites in southern France.


Tabla 1s. Intervenciones de gestión del hábitat realizadas por los cazadores en los tres lugares de estudio del sur de Francia.

	Bourg-Saint-Andéol	Lançon-Provence	Montpeyroux
Openings of scrubland			
Lines (km)	8	12	2
Period of existence	15 years	8 years	3 years
Maintenance frequency	15 ans	3 à 4 ans	3 à 4 ans
Logistic tools	Mechanical shredding + pastoralism	Mechanical shredding	Mechanical shredding
Artisanal feed hoppers			
Unit	0	5	10
Period of existence	-	> 20 years	20 years
Filling frequency (outside of hunting season)	-	Twice to thrice per week	> once per month
Drinking trough			
Unit	2	5	6
Period of existence	> 15 years	> 20 years	> 20 years
Maintenance frequency		Summer: drought event	
Crops for wildlife			
Unit	5	6	1
Area (acre)	0.65	0.75	0.25
Crop rotation	<i>Onobrychis sativa</i> <i>Medicago sativa</i> <i>Triticum aestivum</i> <i>Lolium perenne</i>	<i>Medicago sativa</i> <i>Triticum aestivum</i>	<i>Onobrychis sativa</i> <i>Medicago sativa</i> <i>Triticum aestivum</i> <i>Hordeum vulgare hexastichum</i> × <i>Triticosecale</i>
Land cover (in percent of managed area, 30 ha)			
Openings	20%	16%	4%
Forest trail	2%	1%	1%
Crops	2%	2%	1%
Bare soil	0%	0%	9%
Herb layer	18%	40%	20%
Low scrubland (< 1.5 m)	2%	25%	10%
High scrubland (> 1.5 m)	56%	16%	55%

Table 2s. List of the 17 bird species considered priority species, ordered according to their conservation status in France (IUCN France et al 2016): EN, endangered; VU, vulnerable; NT, near threatened; LC, least concern. (The table also includes the status in annex I of the European Bird Directive and the species main habitat).

Tabla 2s. Lista de las 17 especies de aves consideradas como prioritarias para la conservación ordenadas según el estado de conservación en Francia (IUCN Francia, et al 2016): EN, en peligro de extinción; VU, vulnerable; NT, casi amenazada; LC, preocupación menor. (En la tabla también se indica el estatus de conservación según en el anexo I de la Directiva Europea de Aves y el hábitat principal de la especie).

Species	IUCN French status	Listed in Annex I of the European Birds Directive		Habitat
		No	Yes	
<i>Acrocephalus melanopogon</i>	EN	No		Wetlands
<i>Lanius meridionalis</i>	EN	No		Scrubland, maquis, vine yards
<i>Sylvia conspicillata</i>	EN	No		Mediterranean shrub vegetation
<i>Sylvia undata</i>	EN	No		Mediterranean shrub vegetation
<i>Anthus pratensis</i>	VU	No		Wet grasslands
<i>Carduelis carduelis</i>	VU	No		Heterogenous rural environments
<i>Chloris chloris</i>	VU	No		Heterogenous rural environments
<i>Lanius senator</i>	VU	No		Scrubland, maquis, orchards, dry grasslands
<i>Linaria cannabina</i>	VU	No		Mosaic of agricultural habitat, scrubland, maquis, vineyards
<i>Pyrrhula pyrrhula</i>	VU	No		Temperate forests, groves, parks and gardens
<i>Saxicola rubetra</i>	VU	No		Wet grasslands
<i>Serinus serinus</i>	VU	No		Heterogenous rural environments
<i>Streptopelia turtur</i>	VU	No		Heterogenous rural environments
<i>Coracias garrulus</i>	NT	Yes		Heterogenous rural environments
<i>Lanius collurio</i>	NT	Yes		Semi-open environments with meadows and bushy pastures
<i>Anthus campestris</i>	LC	Yes		Dry grasslands
<i>Caprimulgus europaeus</i>	LC	Yes		Temperate shrubby vegetation

A**B**

	Bourg-Saint-Andéol		Lançon-Provence		Montpeyroux	
Area	MA	UMA	MA	UMA	MA	UMA
Recorded species richness	46	37	53	41	52	43
Estimated species richness	50	41	58	47	56	47
Exhaustivity index (%)	91	90	91	88	93	91

Fig. 1s. A, species accumulation curves for bird species detected at point counts in each managed (MA) and unmanaged (UMA) area of the three study sites in southern France (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux) using the R package 'iNEXT'. B, exhaustivity index for point counts surveys in each area of the three study sites (according to Lauga et Joachim 1987).

Fig. 1s. A, curvas de acumulación de las especies de aves detectadas en los puntos de conteo en cada área gestionada (MA) y no gestionada (UMA) de los tres sitios de estudio en el sur de Francia (BSA, Bourg-Saint-Andéol; LP, Lançon-Provence; MP, Montpeyroux), utilizando el paquete "iNEXT" de R. B, indicador de exhaustividad para los puntos de conteo en cada área de los tres sitios de estudio (según Lauga y Joachim, 1987).